

abberior LIVE 550, Halo, 30 nmol


Item number

LV550-0147-30NMOL

Description

abberior LIVE 550 is a member of a new class of fluorescent dyes for live-cell imaging exclusively offered by abberior. This dye is highly photostable and bright orange fluorescent dye, which can be effectively excited with a 561 nm laser. For STED, a depletion wavelength between 650 - 775 nm is recommended. Due to the unique isomeric structure the favored conformation is stabilized, leading to an enhanced cell permeability of the dye. abberior LIVE 550 can therefore be used in very low nanomolar concentration, reducing background and cytotoxicity effectively, allowing long-term STED and confocal imaging of living cells or tissue. Our abberior LIVE 550 is the ideal partner for abberior LIVE 610 to obtain optimal 2 color live-cell STED and confocal results.

Discover our abberior LIVE 550 Halo, a fluorogenic label that covalently binds to the active site of your HaloTag fusion protein within living cells. This label rapidly binds with exceptional specificity under typical physiological conditions.



Properties

Absorption	λ_{ex} [nm]	551
Extinction Coefficient	ϵ_{max} [M ⁻¹ cm ⁻¹]	75000
Emission	λ_{em} [nm]	573
Quantum Efficiency	η_{fl} [%]	41
STED min.	$\lambda_{\text{STED min}}$ [nm]	650
STED max.	$\lambda_{\text{STED max}}$ [nm]	775
Fluorescence Lifetime	τ_{fl} [ns]	2
Charge	Δq	0
Molecular Weight	MW [g/mol]	636.3
Derivative/Conjugate		Halo

Photophysical properties were measured for carboxylic acid in PBS pH 7.4.

Storage

Each purchase includes 1 vial of lyophilized abberior LIVE 550 Halo, ensuring top-quality performance for live-cell STED and confocal microscopy. For optimal use, we recommend a stock solution of 1 mM in DMF or DMSO. Adjust the working solution concentration between 0.1 μM to 0.5 μM based on the cell type, tissue, and expression rate of your HaloTag fusion protein. HaloTag® labels are shipped freeze-dried in amounts of 30 nmol per vial. Upon arrival, the substances can be stored for up to one year at $-20\text{ }^{\circ}\text{C}$. Shortly before the staining procedure dissolve the label in DMF or DMSO. Once dissolved the stock solutions should be kept at $20\text{ }^{\circ}\text{C}$, protected from light and moisture.

- Note: Depending on solvent quality the shelf-life of the stock solutions might be significantly reduced compared to the substances in their solid form – even if stored at $-20\text{ }^{\circ}\text{C}$.