

abberior LIVE 550, actin, 50 µg


Item number

LV550-0140-50UG

Description

abberior LIVE 550 is a member of a new class of fluorescent dyes for live-cell imaging exclusively offered by abberior. This dye is highly photostable and bright orange fluorescent dye, which can be effectively excited with a 561 nm laser. For STED, a depletion wavelength between 650 - 775 nm is recommended. Due to the unique isomeric structure the favored conformation is stabilized, leading to an enhanced cell permeability of the dye. abberior LIVE 550 can therefore be used in very low nanomolar concentration, reducing background and cytotoxicity effectively, allowing long-term STED and confocal imaging of living cells or tissue. Our abberior LIVE 550 is the ideal partner for abberior LIVE 610 to obtain optimal 2 color live-cell STED and confocal results.

Cell permeable and photostable fluorescent probe used to label actin in fixed and living cells. This highly specific actin probe is based on Jasplakinolide.



Properties

Absorption	λ_{ex} [nm]	551
Extinction Coefficient	ϵ_{max} [M ⁻¹ cm ⁻¹]	75000
Emission	λ_{em} [nm]	573
Quantum Efficiency	η_{fl} [%]	41
STED min.	$\lambda_{\text{STED min}}$ [nm]	650
STED max.	$\lambda_{\text{STED max}}$ [nm]	775
Fluorescence Lifetime	τ_{fl} [ns]	2
Charge	Δq	0
Molecular Weight	MW [g/mol]	1199.5
Derivative/Conjugate	Actin (jasplakinolide)	

Photophysical properties were measured for carboxylic acid in PBS pH 7.4.

Storage

This product contains 1 vial of abberior LIVE 550 actin as lyophilized powder, a stock solution of 1 mM in DMF or DMSO is recommended. Depending on cell type or tissue a working solution of 1 nM to 100 nM is advised. Additionally, a vial containing 1 μ mol verapamil is provided as an imaging reagent. 10 to 20 nmol verapamil in 1 ml staining solution increases the signal intensity significantly. abberior LIVE labels and Verapamil are shipped freeze-dried in amounts of 50 μ g per vial (or 2 nmol in case of evaluation samples) and 1 μ mol, respectively. Upon arrival, the substances can be stored for up to one year at -20 °C. Shortly before the staining procedure dissolve the probe and Verapamil in DMF or DMSO. Once dissolved the stock solutions should be kept at -20 °C, protected from light and moisture.

- Note: Depending on solvent quality the shelf-life of the stock solutions might be significantly reduced compared to the substances in their solid form – even if stored at -20 °C.