


abberior LIVE 550, click, 30 nmol

Item number

LV550-0006-30NMOL

Description

abberior LIVE 550 is a member of a new class of fluorescent dyes for live-cell imaging exclusively offered by abberior. This dye is highly photostable and bright orange fluorescent dye, which can be effectively excited with a 561 nm laser. For STED, a depletion wavelength between 650 - 775 nm is recommended. Due to the unique isomeric structure the favored conformation is stabilized, leading to an enhanced cell permeability of the dye. abberior LIVE 550 can therefore be used in very low nanomolar concentration, reducing background and cytotoxicity effectively, allowing long-term STED and confocal imaging of living cells or tissue. Our abberior LIVE 550 is the ideal partner for abberior LIVE 610 to obtain optimal 2 color live-cell STED and confocal results.



All our bright and photostable abberior LIVE click probes are modified by a tetrazine substrate as a functional group. They react with dienophiles such as trans-cyclooctene (TCO) or bicyclononyne (BCN) derivatives. Along with this bioorthogonal reaction, LIVE click probes are particularly useful in combination with unnatural amino acid (uAA) with a TCO or BCN side chain. The incorporation of uAAs prepares site-selective protein conjugates with a very specific and minimum-sized tag. abberior LIVE click is optimal for super-resolution microscopy within living cells, on cell surfaces, or for in vitro applications.

Properties

Absorption	λ_{ex} [nm]	551
Extinction Coefficient	ϵ_{max} [M ⁻¹ cm ⁻¹]	75000
Emission	λ_{em} [nm]	573
Quantum Efficiency	η_{fl} [%]	41
STED min.	$\lambda_{\text{STED min}}$ [nm]	650
STED max.	$\lambda_{\text{STED max}}$ [nm]	775
Fluorescence Lifetime	τ_{fl} [ns]	2
Charge	Δq	0
Molecular Weight	MW [g/mol]	599.6
Derivative/Conjugate	Click (Tetrazine)	

Photophysical properties were measured for carboxylic acid in PBS pH 7.4.

Storage

This product contains 1 vial of abberior LIVE 550 click as a lyophilized powder, a stock solution of 1 mM in DMF or DMSO is recommended. Depending on cell type, tissue, and expression rate of the unnatural amino acid a working solution of 10 nM to 1 μ M is advised. Upon arrival, the substances can be stored for up to one year at –20 °C. Shortly before the staining procedure dissolve the probe and Verapamil in DMF or DMSO. Once dissolved the stock solutions should be kept at –20 °C, protected from light and moisture.

- Note: Depending on solvent quality the shelf-life of the stock solutions might be significantly reduced compared to the substances in their solid form – even if stored at –20 °C.