

abberior STAR 440L, carboxylic acid, 1 mg


Item number

ST440L-0001-1MG

Description

abberior STAR 440L is a long Stokes-shift dye optimized for STED and confocal microscopy in the blue-green spectral range. It shows a Stokes-shift of approx. 80 nm. As all long Stokes-shift dyes, it shows a broad excitation and emission band: It can be excited between 400 - 470 nm and emits between 490 - 550 nm. For STED microscopy, abberior STAR 440L can be most efficiently used with STED laser wavelengths between 590 - 610 nm. It is therefore well suited for 2 color (STED) imaging together with our abberior STAR GREEN or abberior STAR 488. The hydrophilic properties of abberior STAR 440L conjugates preclude unspecific binding of this dye and thus decrease undesired background fluorescence. Our abberior STAR 440L can substitute dyes like ATTO® 425 or BD Horizon® V500. abberior STAR 440SXP has been renamed abberior STAR 440L. The structure of the dye has not changed. Best results are obtained with freshly prepared samples.

Organic fluorescent dye for microscopy.



Properties

Absorption	λ_{ex} [nm]	432
Extinction Coefficient	ϵ_{max} [M ⁻¹ cm ⁻¹]	33000
Emission	λ_{em} [nm]	511
Quantum Efficiency	η_{fl} [%]	57
STED min.	$\lambda_{\text{STED min}}$ [nm]	590
STED max.	$\lambda_{\text{STED max}}$ [nm]	610
Fluorescence Lifetime	τ_{fl} [ns]	3.7
Correction Factor 260	CF ₂₆₀	0.47
Correction Factor 280	CF ₂₈₀	0.31
Charge	Δq	0
Molecular Weight	MW [g/mol]	528.5
Derivative/Conjugate	Carboxylic acid	

Photophysical properties were measured for carboxylic acid in PBS pH 7.4.

Storage

The product contains a vial of lyophilized powder and is shipped at room temperature. Upon arrival, please store the product at -20 °C. For proper long-term storage up to one year, protect the product from light and moisture.

- Our abberior carboxylic acid derivates come as a freeze-dried powder and should be dissolved in DMF, DMSO, or acetonitrile.