

abberior STAR RED, maleimide, 1 mg


Item number

STRED-0003-1MG

Description

abberior STAR RED is a well-known red dye, named also KK114 in literature. It was developed for STED and confocal imaging in the red spectral region. It shows an exceptional brightness and low background, delivering stunning STED images. abberior STAR RED can be excited with lasers at wavelength between 630 - 650 nm. For STED, a depletion wavelength around 750 - 800 nm is recommended. abberior STAR RED can substitute dyes like ATTO® 647N, Alexa Fluor® 647, or Cy5®. Our abberior STAR RED is the ideal partner for abberior STAR ORANGE or abberior STAR 580 to obtain optimal 2 color STED results. Best results are obtained with freshly prepared samples.

The maleimide is used for conjugating a fluorescent dye to a thiol group (cysteine residues) on a protein or thio-modified molecules. The resulting abberior conjugate will show a bright fluorescence and photostability. abberior offers labeling protocols for labeling of antibodies via thiol-reactive groups.



Properties

Absorption	λ_{ex} [nm]	638
Extinction Coefficient	ϵ_{max} [M ⁻¹ cm ⁻¹]	120000
Emission	λ_{em} [nm]	655
Quantum Efficiency	η_{fl} [%]	90
STED min.	$\lambda_{\text{STED min}}$ [nm]	750
STED max.	$\lambda_{\text{STED max}}$ [nm]	800
Fluorescence Lifetime	τ_{fl} [ns]	3.4
Correction Factor 260	CF ₂₆₀	0.16
Correction Factor 280	CF ₂₈₀	0.32
Charge	Δq	-1
Molecular Weight	MW [g/mol]	1010
Derivative/Conjugate		Maleimide

Photophysical properties were measured for carboxylic acid in PBS pH 7.4.

Storage

The product contains a vial of lyophilized powder and is shipped at room temperature. Upon arrival, please store the product at -20 °C. For proper long-term storage up to one year, protect the product from light and moisture.

- Our abberior maleimides are freeze-dried and should be dissolved only shortly before the coupling reaction. Recommended solvents include amine- and water-free DMF, DMSO, or acetonitrile.